

First photographic record of marbled cat *Pardofelis marmorata* Martin, 1837 (Mammalia, Carnivora, Felidae) in Nepal

Sonam Tashi Lama^{1,2}, James G. Ross¹, Damber Bista², Ang Phuri Sherpa²,
Ganga Ram Regmi³, Madan Krishna Suwal^{3,4}, Pema Sherpa², Janno Weerman⁵,
Shrota Shrestha Lama⁶, Madhuri Thapa⁷, Laxman Prasad Poudyal⁸,
Adrian M. Paterson¹

1 Department of Pest-management and Conservation, Faculty of Agriculture and Life Science, Lincoln University, Ellesmere Junction Road/Springs Road, PO Box 85084, Canterbury, New Zealand **2** Red Panda Network, House No 86, Rani Devi Marga, Ward No 3, Lazimpat, Kathmandu, Nepal **3** Global Primate Network, GPO Box 26288, Kathmandu, Nepal **4** Department of Geography, University of Bergen, Bergen, Norway **5** Rotterdam Zoo, the Netherlands **6** Freelancer, Maharajjung, Kathmandu, Nepal **7** Department of Forests, Ministry of Forests and Environment, Babarmahal, Kathmandu, Nepal **8** Department of National Parks and Wildlife Conservation, Babarmahal, Kathmandu, Nepal

Corresponding author: Sonam Tashi Lama (sonam.tac@gmail.com, Sonam.Lama@RedPandaNetwork.org)

Academic editor: C. Knogge | Received 10 October 2018 | Accepted 10 December 2018 | Published 21 January 2019

<http://zoobank.org/264D80B7-478C-4A65-B5EB-2EA3B015EF6F>

Citation: Lama ST, Ross JG, Bista D, Sherpa AP, Regmi GR, Suwal MK, Sherpa P, Weerman J, Lama SS, Thapa M, Poudyal LP, Paterson AM (2019) First photographic record of marbled cat *Pardofelis marmorata* Martin, 1837 (Mammalia, Carnivora, Felidae) in Nepal. Nature Conservation 32: 19–34. <https://doi.org/10.3897/natureconservation.32.29740>

Abstract

The marbled cat *Pardofelis marmorata* is a Near Threatened small felid. The cat's presence in Nepal is based on an anecdote. A camera trap-based study to assess diversity and abundance of terrestrial mammals in eastern Nepal accumulated 3,014 camera trap days and resulted in 5,176 photographs of 17 medium-large sized mammal species. Amongst them, a marbled cat was captured at a single camera trap station in January 2018. The camera trap-capturing the marbled cat was located in the secondary forest at an altitude of 2,750 m a.s.l., dominated by free-ranging cattle close to a permanent human settlement (1.4 km) and a temporary cattle herding camp (0.4 km). This is the first photographic evidence of a marbled cat in Nepal. In this survey, we also recorded three other felid species: common leopard *Panthera pardus*, Asiatic golden-cat *Catopuma temminckii* and leopard cat *Prionailurus bengalensis*. We recommend detailed year-round camera trap surveys in the mid-hills of eastern Nepal along with research on adaptation of the small

felids to human-dominated areas and assessment of immediate threats for preparing sound conservation management plans of the marbled cat and its sympatric species. Initiation of conservation programmes engaging local *dokpa* (herders) is necessary.

Keywords

Marbled cat, *Pardofelis marmorata*, trail camera, small cats, Eastern Himalayas, Nepal

Introduction

The marbled cat, *Pardofelis marmorata* Martin, 1837, is categorised as ‘Near Threatened’ in the IUCN Red List of Threatened Species (Ross et al. 2016) and nationally data deficient in Nepal (Amin et al. 2018; Jnawali et al. 2011). The marbled cat, listed in CITES Appendix I is a member of the Felidae family along with 14 genera and 42 species of cats (ITIS 2018). This small wild cat is a member of the bay cat lineage along with the Asiatic golden cat *Catopuma temminckii* and bay cat *Catopuma badia*. The marbled cat is similar in colouring to the clouded leopard *Neofelis nebulosa*, with dark blotches, stripes and spots. An arboreal cat with great climbing skills (Mohamed et al. 2009; Sunquist and Sunquist 2017), it is little studied (Sanderson and Watson 2011; Sunquist and Sunquist 2014). The marbled cat is slightly larger and more slender than a domestic cat (Blanford 1888–1891), with a long and bushy tail that is similar in length to the cat’s body. Their sides and back are prominently smudged with irregular, large and dark-fringed markings. Black dots are embellished on legs and underparts and the tail has proximal black spots and distal rings (Sunquist and Sunquist 2017). Out of the two distinct subspecies of marbled cats, *Pardofelis marmorata marmorata* and *Pardofelis marmorata charltonii* (ITIS 2018; Kitchener et al. 2017), the latter is distributed from the foothills of Nepal, through Sikkim, Assam and into upper Myanmar (Pocock 1939). *Pardofelis marmorata charltonii* has a much thicker and longer coat than *P. m. marmorata*, with a rich ochrous-brown colour (Pocock 1939).

The forest-dependent marbled cat is distributed throughout Southeast Asia, from the Himalayan foothills of India and Bhutan to China and then southwards through to Malaysia and Indonesia. The marbled cat has been recorded in Bangladesh (Hance 2016; Khan 2015), Bhutan (Tempa et al. 2013), Brunei Darussalam (Ross et al. 2016), Cambodia (Gray et al. 2014), China (KFBG 2015; Wang and Wang 1986), India (Choudhury 1996), Indonesia (Cheyne and Macdonald 2010), Lao People’s Democratic Republic (Johnson et al. 2009), Malaysia (Azlan and Sharma 2006), Myanmar (Zaw et al. 2014), Thailand (Grassman et al. 2005) and Vietnam (Nowell and Jackson 1996). Although Nepal is noted as the westernmost end of the range in distribution for the marbled cat by the IUCN Red List (Ross et al. 2016), there has been no record of direct sighting or photographic proof of the species in Nepal (Dahal and Dahal 2017).

The distribution map for the marbled cat in Nowell and Jackson (1996) shows Nepal’s Nawalpur, west of Chitwan National Park, as the westernmost record of the species based on a single dead specimen record of the species (Ross et al. 2016). However, the species was not recorded inside the park at that time. The nearest confirmed record

to Nepal is from Khangchendzonga Biosphere Reserve in Sikkim, India and further south in the Manas National Park, in Assam, India. Despite many extensive camera-trapping surveys in Chitwan National Park and other parts of Nepal, there have been no records of the live marbled cat (Dahal and Dahal 2017; Lamichhane et al. 2014; Lamichhane et al. 2016; Yadav et al. 2018). One skin of the species was collected by B. H. Hodgson and presented to the British Museum and labelled as ‘Nepal’ (Pocock 1939). It is assumed that the skin might have been collected during his studies between 1823 and 1843. Blanford (1888–1891) reported the marbled cat as present in Sikkim but not in Nepal. Similarly, a mammal survey report in Nepal by the Bombay Natural History Society found no evidence of the marbled cat in Nepal and speculated that Sikkim may have been the origin of the Hodgson’s specimen (Hinton and Fry 1923). Pocock (1939) declared the claim of Blanford (1888–1891) to be wrong without presenting any reasons. Horsfield (1856) only mentioned ‘hilly region’ not the area as the habitat of marbled cat in his catalogue of the collection of Mammalia from Nepal, Sikkim and Tibet. Pocock (1939) also mentioned that the species was very hard to obtain for collection and seldom received alive by zoos. At that time, the team members of the Bombay Mammal Survey were not able to capture or buy a single animal from the indigenous people in Nepal, Sikkim, Assam and Myanmar. This also indicates the naturally potential rarity of the cat at that time.

The species is primarily found in moist and mixed deciduous-evergreen tropical forest and prefers hilly forests (Duckworth et al. 1999; Grassman et al. 2005; Holden 2001; Nowell and Jackson 1996). The marbled cat is also found in secondary and logged forests, clearings, mangroves and on the periphery of the oil palm plantations (Bernard et al. 2013; Hearn et al. 2016; Ross et al. 2010). The marbled cat’s diet likely consists of squirrels and rats and birds, up to the size of pheasants (Nowell and Jackson 1996; Pocock 1932). This matches what is known from an individual in captivity (Sunquist and Sunquist 2017). Additionally, there is a report of the hunting of a juvenile male Phayre’s leaf monkey by a marbled cat in Thailand (Borries et al. 2014).

The species occurs in isolated and fragmented habitat patches in some part of their distribution range (Ross et al. 2016). The loss of the habitat caused by deforestation is considered as a major threat for the species. Indiscriminate snaring is also common throughout much of its range and is likely a threat to the species (Ross et al. 2016). Body parts of this species were also reported from an illegal market of Yunan, China (Haibin and Kunming 1999). The marbled cat is also hunted for meat (Selvan et al. 2013a) and for socio-cultural rituals by some tribal groups in India (Selvan et al. 2013b). Amongst the numerous confiscations of leopard pelts, there is no record of marbled cat pelts in Nepal (CIB 2018).

Camera traps have been recording and providing new information on distribution and aspects of ecology, in recent times, of small and medium-sized wild felids throughout Southeast Asia, such as the marbled cat (Tempa et al. 2013; Thinley et al. 2015; Wibisono and McCarthy 2010). Camera traps provide absolute evidence that a species is present in an area but are not sufficient to propose conservation action without associating natural densities, habitat quality, quantity, connectivity, land use, hunting pressure etc.

The first ever camera trap photograph of a marbled cat was captured during daylight hours in Huai Kha Khaeng Wildlife Sanctuary in Thailand in 1994 (K. Conforti. pers. comm. as cited in Nowell and Jackson 1996). Similarly, the first photograph of a marbled cat in the wild was taken in Khao Yai Wildlife Sanctuary, Thailand in 1993 (Jackson 1997).

Some areas of Panchthar district (Nepal) are also connected to the protected areas of Sikkim (India) and the habitat area is similar to the habitat in Sikkim, where the occurrence of marbled cat was reported by Balnford (1888–1891). The habitat area in Nepal has much pressure from the anthropogenic activities like free-range cattle grazing, fuel wood and fodder collection.

Methodology

A camera-trapping study was conducted in non-protected forest areas of the Kangchenjunga landscape, bordering India on the eastern side, connecting the Singhalila National Park, Barsey Rhododendron Sanctuary and Khangchendzonga Biosphere Reserve, in far eastern Nepal (26°59'6.159"–27°25'57.969" N and 87°51'24.084"–88°6'34.282" E) (Figure 1). The elevation range of the study area lies between 2,000 to 4,355 m above sea level (m a.s.l.), covering about 520 km². The study area has been proposed as a conservation corridor to connect Nepal's Kangchenjunga Conservation Area (KCA) to these Indian conservation areas (Chettri et al. 2007). Monsoonal rain decreases from the beginning of October and the colder days increase in frequency, with the leaves of the deciduous hardwoods starting to fall by the end of November. This study area has been identified as a last remaining tract of Eastern Himalayan broadleaf forest, a major habitat for the endangered red panda (Williams 2003).

Dominant forest species and the composition of forest species changes with elevation, for instance *Quercus* sp. in lower temperate forest (1700–2400 m a.s.l.), *Quercus* sp., *Rhododendron* sp., and *Acer* sp., *Pinus* sp. in upper temperate forest (2400–2800 m a.s.l.), *Rhododendron* sp., *Betula* sp. and *Acer* sp. in subalpine forest (around 3000 m a.s.l.) and shrubby rhododendron species and junipers in the alpine zone (Dombremez 1976, as cited in Jackson 1994). Free range grazing has been in practice in the region for more than 170 years (Hooker 1889), is prevalent above 2,500 m a.s.l., impacting the forest area with over-grazing and has been listed as a major threat for biodiversity conservation (Shrestha et al. 2008).

A camera trap study was conducted to inventory the medium- to large-sized mammalian fauna in the region between December 2017 and June 2018, encompassing two seasons; winter (December 2017–February 2018) and spring (March–June 2018). The whole study area (Figure 1) was considered as a single block. The area above 2,000 m altitude was first selected by geoprocessing in ArcGIS 10.2. Then, we imposed uniform grid cells of 2 km × 2 km throughout the selected area. This resulted in 211 grids, of which 60 grids were sampled based on habitat type and accessibility. During field sampling, we omitted private forest areas and difficult rocky mountain




Figure 1. A map of the study area in Southern Kangchenjunga Landscape in eastern Nepal showing sampled grids in blue and location of the marbled cat detection.

cliffs and steep-mountain slopes because of inaccessibility. Within a selected grid, active trails and major areas of interest to wildlife, containing water sources, denning areas, major passes along mountain ridges and mineral licks, were chosen for the installation site of the camera traps.

We used 63 passive infrared camera trap units (51 Bushnell TROPHY Cam HD camera trap units from Bushnell Outdoor Products, USA and 12 units of Strike Force Elite camera traps, Model BTC-5HDE from Browning, Australia). The location and altitude of each site were recorded using a Garmin eTrex 10 and Rino120 handheld GPS.

On average, the cameras were left in the forests for 28 trap days. The minimum trap days for a camera was 14 days while 78 days was the maximum. The camera traps were installed at 107 locations. Cameras were secured to available tree trunks. The sensor height of the cameras installed above ground level was 32 cm on average. Every 24 hours was considered as one camera trap day for this study. The photographs were sorted manually and also with the help of Microsoft Access-based Camera Base 1.7 (Tobler 2015). The species from the photographs were identified with the help of the references from Sanderson and Watson (2011), Johnsingh and Manjrekar (2012) and Menon (2014), as well as consultation with experts and Forest Guardians from local villages who assisted with installing the camera traps. Photographs that were taken more than 30 minutes apart from each other were considered as independent events (Di Bitetti et al. 2006; Silver et al. 2004).

Results and discussions

Overall, there were 3,014 trap days (1,402 in winter and 1,612 in spring) from the 107 camera stations (52 in winter and 55 in spring) that collected 5,176 photographs of 17 medium-large sized mammal species. Of interest were three photos of a small felid (two whole body and one tail-only photo from its right flank) in a single event from a location ID W33 (27°20'2.22"N, 88°0'55.08"E; datum WGS84), (elevation: 2,750 m a.s.l.; aspect: west; slope: steep, details on Table 1) an dusk at 17:44 hr on 21 January 2018 (Figure 2). During the successive spring season (March-June), we surveyed the area where the marbled cat was detected intensively and systematically with seven monitoring stations. This resulted in 233 more camera days but with no more images of the marbled cat. One of the cameras was installed in previously trapped locations for a further 35 days.

The cat recorded location was near Tham Chok (a temporary shed for *dokpa*) from Phalaicha village in Panchthar District. The photos of the small felid were markedly different from other small cat images captured by other cameras in the area. A long, fluffy and erect tail, black edged large blotches on a flank and solid spots on the limbs were consistent with a marbled cat (confirmed by Igor Khorozyan, Jim Sanderson, Karan Shah, Kashmira Kakati, Rinjan Shrestha and Yadav Ghimirey). The camera trap images also matched (Figure 2) with published photos of the species (Dhendup et al. 2016; Nowell and Jackson 1996; Sethy et al. 2017). The marbled cat images clearly differed from the other felid species observed in this study, i.e. Asiatic golden cat, common leopard and leopard cat. The confirmation of the presence of the marbled cat at this site supports C. MacDougal's suggestion (Nowell and Jackson 1996) about the presence of the marbled cat in Nepal despite the lack of clear evidence about the origin of the cat's skin.

Table 1. Camera trap location details where the marbled cat was photographed in Southern Kangchenjunga region, eastern Nepal.

| Parameter | Value |
|---|--|
| Location ID/Name of Location | W33/Tham Chok |
| Long/Lat | 27°20'2.22"N, 88°0'55.08"E |
| Elevation | 2,750 m |
| Total number of photos | 3 |
| No. of Independent events | 1 |
| Duration of camera placement | 7–22 Jan 2018 (15 days) |
| Date, time & temperature of marbled cat recorded photo | 21-01-2018, 17h 44m & 5°C |
| Habitat type | Upper-temperate mixed-broadleaved forest |
| Site aspect | West |
| Slope | Steep |
| Camera sensor height from ground | 50 cm |
| Distance to nearest human settlement | 1.4 km |
| Distance to nearest cattle herding station | 0.4 km |
| Distance to forest edge | 320 m |
| Distance to permanent water source | 1 km |
| Distance to nearest forest in Sikkim, India | 4 km |
| Distance to nearest trail for human & livestock | 50 m |
| Other mammal species captured at the same camera | Orange-bellied Himalayan squirrel & Northern red muntjac |
| Bird species captured in the same camera | Kalij-pheasant |
| Species captured by the camera trap in same location in spring season | Leopard cat, Red muntjac, Yellow-throated marten and Wild boar |

The habitat of the Tham Chok area is upper-temperate broadleaved forest (Figure 3). The vegetation in the area is dominated by *Lithocarpus pachyphylla*, *Castanopsis hystrix*, *Rhododendron grande*, *Taxus wallichiana*, *Lyonia ovalifolia* and *Alnus nepalensis* with thickets of *Arundinaria* sp. The recorded site is close to the Iwaa River, which originates from Timbung Lake at 4,200 m a.s.l.. The ridge is just above the confluence of Iwaa and Barne Rivers, which stretch towards the border with India, approximately 4 kilometres (km) away. The area is a summer grazing pasture for cows and goats and also a winter grazing pasture for dzomu (domestic yak-cow hybrids), supporting around 150 livestock over the winter. The eastern part of the Tham Chok is a largely rocky area with dwarf trees, probably recovering from a historical forest fire. A large area of primary forest in the Iwaa valley was burnt by a forest fire, presumably caused by incidental poachers, approximately 3.5 decades ago (pers. comm. K. Bhandari 2018). That forest fire caused huge damage to the local fauna and flora at that time. The habitat area is crisscrossed by human and livestock trails. A human trail was 50 m away from the camera trap station that recorded the marbled cat. The elevation record of the species in this study at 2,750 m a.s.l. is higher than the elevation limit of the species for Nepal (< 2,500 m a.s.l.) presented by Baral and Shah (2008). Despite a heavy presence of humans and livestock in the area, there is some pristine primary forest dominated by *Lithocarpus pachyphylla*, *Castanopsis hystrix*, and *Rhododendron* sp., in



Figure 2. An image of the first marbled cat in Nepal from the camera trap (Sonam Tashi Lama-Lincoln University/Red Panda Network).

the eastern portion of the area, approximately 2.5 km away. That primary forest could be a refuge habitat for many wildlife species, including small and medium-sized felids.

The other mammal and bird species captured at the same camera trap were the northern red muntjac *Muntiacus vaginalis*, orange-bellied Himalayan squirrel *Dremomys lokriah* and the Kalij pheasant *Lophura leucomelanos*. Similarly, leopard cat, yellow-throated marten *Martes flavigula* and wild boar *Sus scrofa* were recorded in the spring season. The photo-capture rate for that camera was highest for the muntjac followed by the squirrel in winter. The main prey present in the area, the orange-bellied squirrels (Figure 4) (Nowell and Jackson 1996) were also found to be strongly diurnal of habit in this study. Our images also support that marbled cat was primarily diurnal in their habit (Figure 2) as many other records around the world. A nearby camera trap, approximately 1 km away, only captured the photographs of dzomus, indicating the pressure of livestock in the area.

Marbled cat appear to have a very low capture rate throughout their range. The reason behind the low rate of the capture of the species could be due to the rareness of the species itself (Nowell and Jackson 1996; Azlan and Sharma 2006; Cheyne and Macdonald 2010). Failure to record the marbled cat during the spring season, despite intensive sampling of locations, may also indicate the rareness of the species in our study area or that we missed them because all of our cameras were based on the



Figure 3. The landscape and forests of the marbled cat capture area in Iwaa valley. The cat was photographed at far right (marked with red oval). (Sonam Tashi Lama-Lincoln University/Red Panda Network).

ground and they can be arboreal in nature (Sunquist and Sunquist 2017). Azlan and Sharma (2006) also reported a very low abundance of marbled cat (only one photo) and suggested that the cat could be rare and/or are mostly arboreal. Cheyne and Macdonald (2010), in Sabangau peat-swamp forest, Indonesian Borneo, did succeed in trapping a single image of this cat over 15 months, which further suggests the rarity of this species in the natural habitat. Camera-trap images of the marbled cat and the leopard cat were shown to nine *dokpa* in the first week of October 2018 and asked if they had sighted the cat in their area. Four of *dokpa* reported sighting the marbled cat, but all of them named it as a *chari bagh* (leopard cat). They understood them as a poultry pest that sometimes preys on goats. They had knowledge of the use of marbled cat skin and bones.

In a study by Hearn et al. (2016), the marbled cat was recorded from selectively logged area to primary forests but was not recorded inside the plantation forest. Their study estimated the density of the cat as higher than the Sunda clouded leopard *Neofelis diardi* and quite similar to the estimate of the leopard cat. In our study, the record of the leopard cat was relatively high.

The records of a marbled cat skull by Pocock (1939) from adjoining Darjeeling, India and one skin record originating from Sikkim, indicate that the cats were present in the region decades ago. This was also supported by the recent camera-trapped marbled



Figure 4. The Orange-bellied Himalayan squirrel *Dremomys lokriah*, a probable primary prey of marbled cat, is abundant in the study area. This individual was captured by the same camera trap (Sonam Tashi Lama-Lincoln University/Red Panda Network).

cat in Sikkim, adjoining and contiguous to our study area (pers. comm. P. S. Ghose 2018). This is consistent with the specimen collected by Hodgson from Nepal and the listing this species in the Nepal list by Horsfield (1856).

Hodgson was not allowed to travel outside of Kathmandu Valley during his stay in Nepal but he recruited a team of hunters and trappers to collect specimens for him (Datta and Inskipp 2004). He also did the same during his time (1845–1858) in Darjeeling, India (Hunter 1896) and his four assistants (bird and animal shooters, collector and taxidermist) travelled to eastern Nepal from October through to December 1848 with Sir Josheph Dalton Hooker (Hooker 1889). Due to the requests from the Lamas of Buddhists monasteries not to shoot or even fish in their surroundings (Hunter 1896), this might have limited the collection. On the other hand, some local ethnic people in eastern Nepal worked as licensed hunters for four generations to British India to hunt and trap birds and mammals for collections (pers. comm. S. B. Rai 2017).

The forest area, where the marbled cat was detected, has been naturally restored as a secondary forest after a forest fire that occurred nearly 3.5 decades ago. The slash-and-burn agriculture system was practised in the adjoining areas for many years until around 1998 (pers. comm. K. Bhandari 2018). The ongoing restoration of the secondary forests supports healthy biodiversity, including the top predators and provides

opportunities for combining forest regeneration and sustainable rural livelihoods for local people (Chazdon 2008). After banning grazing in the forest and removing mobile pastoralism in Sikkim in 1998 and Singhalila in 1992, the pressure of livestock has concentrated into the study area (Thomas 2014). The current number of livestock (cow, buffalo, yak, sheep and goat) in surrounding villages (Phalaicha and Kalikhola only) is 6,472 (MoLD 2017), of which at least 50% of them are grazing freely in nearby forests.

Small cats have been little studied in Nepal (Thapa 2014). The conflict between humans and marbled cat throughout its range appears to be low (Inskipp and Zimmermann 2009). It is not clear whether the low level of knowledge about the marbled cat throughout its range is due to rarity or its solitary and cryptic nature as a species (Sunquist and Sunquist 2017). In this context of limited knowledge on its ecology, distribution and status on a global level (Ross et al. 2016), we stress further research for gathering baseline data and appeal for conservation programmes outside the protected areas of Nepal. We expect to find the presence of the marbled cat in KCA, Makalu Barun National Park and most of the forested areas in the mid-hills of Nepal as the habitat and climates are similar to where we photographed the cat in eastern Nepal. As the coverage of forested areas by a camera-trapping survey in western Nepal is very low and limited to the protected areas, we cannot definitively conclude its absence from that region.

Based on a single presence location recorded in this study, it would be impractical to conclude anything about habitat characteristics for the marbled cat in Nepal. Limited information on population ecology, interactions with other sympatric felids, as well as how the species responds with the growing anthropogenic interactions in human-dominated landscapes, does not allow for an effective conservation plan. Nevertheless, the record of this species within a human-dominated landscape underpins the need for wildlife-friendly conservation measures to secure human-wildlife coexistence. Potential habitat identification following habitat zonation measures will help limit human activities outside the core habitat thus minimising the threat to some extent. However, an in-depth study on habitat characteristics, diet, distribution, movement and activity pattern will further guide in formulating an effective conservation plan targeting this species.

Acknowledgements

We are grateful to the Ministry of Forests and Environment, Department of Forests and Soil Conservation, for granting permission (72-2074/2075) to conduct this camera-trapping study. This research was funded by Rotterdam Zoo, Red Panda Network, Lincoln University and Wildlife Conservation Network. We thank all the forest guardians, especially Jiwan Rai and Sunil Rai (Deep Jyoti Youth Club) for their fieldwork. We thank Global Primate Network Nepal for providing camera traps which were supported to GRR by Mohamed bin Zayed Species Conservation Fund and Small Wild Cat Conservation Foundation. STL dedicates this paper to late Mingma Norbu Sherpa for his contributions to the conservation of nature in the Eastern Himalaya and is also

indebted to Lincoln University, Russell E. Train Education for Nature/WWF and The Greater Himalayas Foundation for Mingma Norbu Sherpa Memorial scholarship. We also greatly appreciate the helpful comments from Igor Khorozyan on a previous version of a manuscript, reviewers J. Sanderson and Christoph Knogge for their constructive comments and Tshiring Lama and Tenzing Y. Bhutia for interviewing *dokpas* in alpine pastures during summer of 2018.

References

- Amin R, Baral HS, Lamichhane BR, Poudyal LP, Lee S, Jnawali SR, Acharya KP, Upadhyaya GP, Pandey MB, Shrestha R, Joshi D, Griffiths J, Khatriwada AP, Subedi N (2018) The status of Nepal's mammals. *Journal of Threatened Taxa* 10(3): 11361–11378. <https://doi.org/10.11609/jott.3712.10.3.11361-11378>
- Azlan JM, Sharma DS (2006) The diversity and activity patterns of wild felids in a secondary forest in Peninsular Malaysia. *Oryx* 40(1): 36–41. <https://doi.org/10.1017/S0030605306000147>
- Baral HS, Shah KB (2008) *Wild Mammals of Nepal*. Himalayan Nature Kathmandu, Nepal, 188 pp.
- Bernard H, Brodie JF, Giordano AJ, Ahmad AH, Sinun W (2013) Bornean felids in and around the Imbak Canyon conservation area, Sabah, Malaysia. *Cat News* 58: 44–46. <http://eprints.ums.edu.my/id/eprint/19056>
- Bhandari K (2018) Interview (recorded) by the first author on 11 January 2018.
- Blanford WT (1888–1891) *Fauna of British India, including Ceylon and Burma – Mammalia*. Taylor and Francis, Red Lion Court, Fleet, Street, 617 pp.
- Borries C, Primeau ZM, Ossi-Lupo K, Dtubpraserit S, Koenig A (2014) Possible predation attempt by a marbled cat on a juvenile Phayre's leaf monkey. *Raffles Bulletin of Zoology* 62: 561–565.
- Chazdon RL (2008) Beyond deforestation: Restoring forests and ecosystem services on degraded lands. *Science* 320(5882): 1458–1460. <https://doi.org/10.1126/science.1155365>
- Chettri N, Sharma E, Shakya B, Bajracharya B (2007) Developing forested conservation corridors in the Kangchenjunga landscape, eastern Himalaya. *Mountain Research and Development* 27(3): 211–214. <https://doi.org/10.1659/mrd.0923>
- Cheyne SM, Macdonald DW (2010) Marbled cat in Sabangau peat-swamp forest, Indonesian Borneo. *Cat News* 52: 11–12. http://www.outrop.com/uploads/7/2/4/9/7249041/cheyne_macdonald_2010_marbled_cat_news.pdf
- Choudhury A (1996) The marbled cat *Felis marmorata* Martin in Assam—some recent records. *Journal of the Bombay Natural History Society* 93: 583–583.
- CIB (2018) Press releases Central Investigation Bureau Nepal Police Kathmandu. <https://cib.nepalpolice.gov.np/index.php/news/press-release> [Retrieved October 4, 2018]
- Dahal BV, Dahal S (2017) Does the marbled cat exist in Nepal? *Cat News* 66: 31–33.

- Datta A, Inskipp C (2004) Zoology... amuses me much. The Origins of Himalayan Studies. Routledge Curzon, London and New York, 158–177. <https://www.taylorfrancis.com/books/e/9781134383641/chapters/10.4324%2F9780203480359-15>
- Dhendup T, Tempa T, Tenzin U (2016) Clouded leopard co-exists with six other felids in Royal Manas National Park, Bhutan. *Cat News* 63: 24–27.
- Di Bitetti MS, Paviolo A, De Angelo C (2006) Density, habitat use and activity patterns of ocelots (*Leopardus pardalis*) in the Atlantic Forest of Misiones, Argentina. *Journal of Zoology* 270: 153–163. <https://doi.org/10.1111/j.1469-7998.2006.00102.x>
- Duckworth W, Salter RE, Khounbolin K (1999) Wildlife in Lao PDR: 1999 status report. World Conservation Union, Lao PDR, 1–275. https://cmsdata.iucn.org/sites/dev/files/import/downloads/lao_wildlife_status_report_1999.pdf
- Ghose PS (2018) Email communication and phone interview by the first author on 31 August and 10 September 2018.
- Grassman Jr LI, Tewes ME, Silvy NJ, Kreetiyutanont K (2005) Ecology of three sympatric felids in a mixed evergreen forest in north-central Thailand. *Journal of Mammalogy* 86(1): 29–38. [https://doi.org/10.1644/1545-1542\(2005\)086<0029:EOTSFI>2.0.CO;2](https://doi.org/10.1644/1545-1542(2005)086<0029:EOTSFI>2.0.CO;2)
- Gray TNE, Phan C, Chanrattanak P, Sovanna P (2014) The status of jungle cat and sympatric small cats in Cambodia's Eastern Plains. *Cat News* 8: 19–23.
- Haibin W, Kunming G (1999) Wildlife trade in Yunnan Province, China, at the border with Vietnam. *Traffic Bulletin* 18: 21–30.
- Hance J (2016) Tiger country? Scientists uncover wild surprises in tribal Bangladesh. *The Guardian*. Guardian News and Media Limited London, UK. <https://www.theguardian.com/environment/radical-conservation/2016/mar/01/tiger-country-scientists-uncover-wild-surprises-in-tribal-bangladesh>
- Hearn AJ, Ross J, Bernard H, Bakar SA, Hunter LT, Macdonald DW (2016) The first estimates of marbled cat *Pardofelis marmorata* population density from Bornean primary and selectively logged forest. *PLoS One* 11(3): e0151046. <https://doi.org/10.1371/journal.pone.0151046>
- Hinton MAC, Fry TB (1923) Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Report No. 37: Nepal. *Journal of the Bombay Natural History Society* 29: 399–428.
- Holden J (2001) Small cats in Kerinci Seblat National Park, Sumatra, Indonesia. *Cat News* 35: 11–14.
- Hooker JD (1889) Himalayan Journals; or Notes of a Naturalist. In: Bettany GT (Ed.) Bengal, the Sikkim and Nepal Himalayas, the Khasia Mountains, &c. Chapter VIII. Ward, Lock and Co., LTD. London, 574.
- Horsfield T (1856) Catalogue of a collection of Mammalia from Nepal, Sikkim, and Tibet, presented to the Hon. East India Company by B. H. Hodgson, Esq., in 1853. *Proceedings of the Zoological Society of London Part XXIV*.
- Hunter WW (1896) Life of Brian Houghton Hodgson: British Resident at the Court of Nepal J. Murray, Albemarle Street, London, 390 pp.

- Inskipp C, Zimmermann A (2009) Human-felid conflict: A review of patterns and priorities worldwide. *Oryx* 43(01): 18–34. <https://doi.org/10.1017/S003060530899030X>
- ITIS (2018) Integrated Taxonomic Information System on-line database. <https://www.itis.gov/> [Retrieved November 27, 2018]
- Jackson JK (1994) Manual of afforestation in Nepal. Volume 1. Forest Research and Survey Centre, Ministry of Forests and Soil Conservation. Kathmandu, Nepal, 327 pp.
- Jackson P (1997) The status of cats in the wild. *International Zoo Yearbook* 35(1): 17–27. <https://doi.org/10.1111/j.1748-1090.1997.tb01185.x>
- Jnawali S, Baral H, Lee S, Acharya K, Upadhyay G, Pandey M, Shrestha R, Joshi D, Laminchane B, Griffiths J [Eds] (2011) The Status of Nepal Mammals: The National Red List Series, Department of National Parks and Wildlife Conservation Kathmandu, Nepal, 266 pp.
- Johnsingh AJT, Manjrekar NE (2012) Mammals of South Asia Volume One. Universities Press (India) Private Limited, Hyderabad, India, 614 pp.
- Johnson A, Vongkhamheng C, Saithongdam T (2009) The diversity, status and conservation of small carnivores in a montane tropical forest in northern Laos. *Oryx* 43(04): 626–633. <https://doi.org/10.1017/S0030605309990238>
- KFBG (2015) Elusive Marbled Cat rediscovered in China's Yunnan Province after 30 years. Kadoorie Farm and Botanic Garden (KFBG) Corporation, Lam Kam Road, Tai Po, New Territories, Hong Kong. <https://www.kfbg.org/eng/news/Marbled-Cat.aspx>
- Khan M (2015) First confirmed record of marbled cat in Bangladesh. *Cat News* 62: 17–17.
- Kitchener AC, Breitenmoser-Würsten C, Eizirik E, Gentry A, Werdelin L, Wilting A, Yamaguchi N, Johnson WE (2017) A revised taxonomy of the Felidae: The final report of the Cat Classification Task Force of the IUCN Cat Specialist Group. *Cat News* 11: 1–80. https://repository.si.edu/bitstream/handle/10088/32616/A_revised_Felidae_Taxonomy_CatNews.pdf
- Lamichhane BR, Dhakal M, Subedi N, Pokheral CP (2014) Clouded leopard co-exist with other five felids in Chitwan National Park, Nepal. *Cat News* 61: 30–32.
- Lamichhane BR, Kadariya R, Subeidi N, Dhakal BK, Dhakal M, Thapa K, Acharya KP (2016) Rusty-spotted cat: 12th cat species discovered in Western Terai of Nepal. *Cat News* 64: 30–33.
- Menon V (2014) Indian Mammals: A Field Guide. Hachette India, 528 pp.
- Mohamed A, Samejima H, Wilting A (2009) Records of five Bornean cat species from Deramakot Forest Reserve in Sabah, Malaysia. *Cat News* 51: 12–15. http://www.cloudedleopard.org/Documents/Mohamed_et_al_Cat_News_51.pdf
- MoLD (2017) Livestock Statistics of Nepal. In: Planning MaED (Ed.) Government of Nepal, Ministry of Livestock Development, Singhadurbar, Kathmandu, Nepal, 1–170.
- Nowell K, Jackson P (Eds) (1996) Wild Cats: Status Survey and Conservation Action Plan. IUCN Gland, 382 pp.
- Pocock RI (1932) The marbled cat (*Pardofelis marmorata*) and some other oriental species, with the definition of a new genus of the Felidae. *Proceedings of the Zoological Society of London*. Wiley Online Library, 741–766. <https://doi.org/10.1111/j.1096-3642.1932.tb01095.x>

- Pocock RI (1939) The Fauna and British India including Ceylon and Burma. Taylor and Francis Ltd., London, E.C. 4, 463 pp.
- Rai SB (2017) Interview (recorded) by the first author on 31 December 2017.
- Ross J, Hearn AJ, Bernard H, Secoy K, Macdonald D (2010) A Framework for a Wild Cat Action Plan for Sabah. Global Canopy Programme. Oxford, 49 pp.
- Ross J, Brodie J, Cheyne S, Datta A, Hearn A, Loken B, Lynam A, McCarthy J, Phan C, Rasphone A, Singh P, Wilting A (2016) *Pardofelis marmorata*. The IUCN Red List of Threatened Species 2016: e.T16218A97164299.
- Sanderson JG, Watson P (2011) Small wild cats: The animal answer guide. John Hopkins University Press, Baltimore, 144 pp.
- Selvan KM, Veeraswami GG, Habib B, Lyngdoh S (2013a) Losing threatened and rare wildlife to hunting in Ziro Valley, Arunachal Pradesh, India. *Current Science* 104: 1492–1495.
- Selvan KM, Gopi GV, Habib B, Lyngdoh S (2013b) Hunting record of endangered marbled cat *Pardofelis marmorata* in the Ziro valley of Lower Subansiri, Arunachal Pradesh, India. *Journal of Threatened Taxa* 5(1): 3583–3584. <https://doi.org/10.11609/JoTT.o3208.100>
- Sethy J, Gouda S, Chauhan NS (2017) Photographic evidence of marbled cat from Dampa Tiger Reserve, Mizoram, India. *Cat News* 65: 26–27.
- Shrestha KK, Kunwar RM, Dhamala MK, Humagain K (2008) Conservation of Plant Resources in Kanchenjunga-Singalila Ridge, Eastern Nepal. *Nepal Journal of Plant Sciences* 2: 62–68.
- Silver SC, Ostro LE, Marsh LK, Maffei L, Noss AJ, Kelly MJ, Wallace RB, Gómez H, Ayala G (2004) The use of camera traps for estimating jaguar *Panthera onca* abundance and density using capture/recapture analysis. *Oryx* 38(02): 148–154. <https://doi.org/10.1017/S0030605304000286>
- Sunquist F, Sunquist M (2014) The Wild Cat Book: Everything you ever wanted to know about cats. University of Chicago Press, 280 pp. <https://doi.org/10.7208/chicago/9780226145761.001.0001>
- Sunquist M, Sunquist F (2017) Wild Cats of the World. University of Chicago Press, Chicago, 462 pp.
- Tempa T, Hebblewhite M, Mills LS, Wangchuk TR, Norbu N, Wangchuk T, Nidup T, Dendup P, Wangchuk D, Wangdi Y, Dorji T (2013) Royal Manas National Park, Bhutan: A hot spot for wild felids. *Oryx* 47(02): 207–210. <https://doi.org/10.1017/S0030605312001317>
- Thapa S (2014) A checklist of mammals of Nepal. *Journal of Threatened Taxa* 6(8): 6061–6072. <https://doi.org/10.11609/JoTT.o3511.6061-72>
- Thinley P, Morreale S, Curtis P, Lassoie J (2015) Diversity, occupancy, and spatio-temporal occurrences of mammalian predators in Bhutan's Jigme Dorji National Park. *Bhutan Journal of Natural Resources & Development* 2: 19–27.
- Thomas S (2014) A Tragedy on the Commons. *Current Conservation* 5.3. <https://www.currentconservation.org/a-tragedy-on-the-commons/>
- Tobler M (2015) Camera Base 1.7 The Botanical Research Institute of Texas. Atrium Biodiversity Information System. <http://www.atrrium-biodiversity.org/tools/camerabase/>

- Wang Z, Wang S (1986) Distribution and recent status of the Felidae in China. In: Miller SD, Everett DD (Eds) *Cats of the World: Biology, Conservation and Management*. National Wildlife Federation, Washington DC, 201–210.
- Wibisono HT, McCarthy J (2010) Melanistic marbled cat from Bukit Barisan Selatan National Park, Sumatra, Indonesia. *Cat News* 52: 9–10.
- Williams BH (2003) Red panda in eastern Nepal: how do they fit into ecoregional conservation of the eastern Himalaya. *Conservation Biology in Asia* 16: 236–250. https://conbio.org/images/content_groups/Asia/Chapter%2016_Williams%20Red%20panda%20Nepal.pdf
- Yadav SKY, Lamichhane BR, Subedi N, Dhakal M, Thapa RK, Poudyal LP, Dahal BR (2018) Fishing cat camera trapped in Babai Valley of Bardia National Park, Nepal. *Cat News* 67: 31–33. <http://www.ntnc.org.np/publication/fishing-cat-camera-trapped-babai-valley-bardia-national-park-nepal>
- Zaw T, Myint T, Htun S, Po SHT, Latt KT, Maung M, Lynam A (2014) Status and distribution of smaller cats in Myanmar. *Cat News* 8: 24–30. <https://agriknowledge.affrc.go.jp/RN/3020062782>