

Uncertainty and the design of in-situ biodiversity-monitoring programs

William E. Magnusson¹

¹ Instituto Nacional de Ciência Tecnologia - Centro de Estudos Integrados da Biodiversidade Amazônica (INCT-CENBAM), Instituto Nacional de Pesquisas da Amazônia (INPA), CP 2223, 69080-971, Manaus AM, Brasil

Corresponding author: William E. Magnusson (bill@inpa.gov.br)

Academic editor: P.-Y. Henry | Received 5 July 2013 | Accepted 6 March 2014 | Published 9 October 2014

<http://zoobank.org/271EE398-5067-4A4D-B4E0-492A45B1C6AA>

Citation: Magnusson WE (2014) Uncertainty and the design of in-situ biodiversity-monitoring programs. Nature Conservation 8: 77–94. doi: 10.3897/natureconservation.8.5929

Abstract

There are many techniques to deal with uncertainty when modeling data. However, there are many forms of uncertainty that cannot be dealt with mathematically that have to be taken into account when designing a biodiversity monitoring system. Some of these can be minimized by careful planning and quality control, but others have to be investigated during monitoring, and the scale and methods adjusted when necessary to meet objectives. Sources of uncertainty include uncertainty about stakeholders, who will monitor, what to sample, where to sample, causal relationships, species identifications, detectability, distributions, relationships with remote sensing, biotic concordance, complementarity, validity of stratification, and data quality and management. Failure to take into account any of these sources of uncertainty about how the data will be used can make monitoring nothing more than monitoring for the sake of monitoring, and I make recommendations as to how to reduce uncertainties. Some form of standardization is necessary, despite the multiple sources of uncertainty, and experience from RAPELD and other monitoring schemes indicates that spatial standardization is viable and helps reduce many sources of uncertainty.

Keywords

Biodiversity, stakeholder, sampling, identification, detectability, distribution, remote sensing, concordance, complementarity, stratification, data quality

Introduction

There are many sources of uncertainty in scientific research, some of which can be modeled mathematically, but some sources of uncertainty are considered non-probabilistic, and the best way to deal with these is controversial (e.g. Sniedovich 2014). Some uncertainties can be reduced by careful quality control during data collection, but most researchers attempt to reduce uncertainty by statistical analysis, usually through model-based inferences (Gitzen et al. 2012, Thornton et al. 2014). There is an enormous literature on model-based inference (e.g. Anderson 2008), but even the most complex of these models, with many alternative hypothesis, relate to only very simple systems with limited inputs and outputs. They are good for evaluating sources of uncertainty that can be modeled by probability distributions, and in essence are usually just attempts to parameterize a given generic model. For example, Anderson (2008) used Caley's and Hone's (2002) study of tuberculosis transmission in ferrets as an example of a multi-hypothesis study. However, all of the hypotheses presented related to ways that ferrets could contract bovine tuberculosis. These are very interesting questions, but a biologist interested in biodiversity questions might have to deal with uncertainty as to whether bovine tuberculosis is better avoided or treated, whether management costs outweigh the costs of no action, whether other species are more important than ferrets in transmission, whether control measures might be considered inhumane, whether climate change or changes in markets might make the question irrelevant, and many other sources of uncertainty that are difficult to put into a probabilistic framework. While it is reasonable to ignore such concerns in a short-term study designed to find a solution to an immediate pressing problem, researchers interested in conserving biodiversity over the next century do not have the luxury of being able to use such a focused and short-term approach (Haila et al. 2014).

Ecologists are generally most worried about uncertainty in relation to their field of research. Taxonomists worry about the correctness of identifications, modelers worry about the accuracy of parameter estimates in their models, sociologists are concerned with uncertainties about the contributions of different stakeholders, geneticists try to reduce uncertainty about gene flow, etc. However, biodiversity managers have to deal with all sources of uncertainty simultaneously, and the importance of different forms of uncertainty will vary depending on the objectives of management. I have therefore adopted a very broad concept of uncertainty, and make recommendations as to how they can be reduced or quantified by planning during implementation of field infrastructure and quality control during data collection.

Here we will describe some of the sorts of uncertainty that we had to take into account when developing the RAPELD system of biodiversity monitoring (Magnusson et al. 2005). We will use that system to illustrate the issues, but the same considerations are applicable to any *in-situ* biodiversity monitoring system. The RAPELD system is a standardized monitoring scheme developed to allow integrated analyses of biodiversity data collected in rapid assessments (RAP) and long-term (LTER [PELD in Portuguese]) studies (Costa and Magnusson 2010; Magnusson et al. 2013). Sampling

in the RAPELD system is based on spatially standardized transects and plots, reducing uncertainty about spatial interpolations and extrapolations. The system can be adapted to different sized areas of interest, but, because it is modular, statistically valid comparisons can be made between studies that originally used different combinations of spatial modules.

The basic sampling unit in the RAPELD system is a 1 km transect combined with one or more permanent plots that are usually 250 m long in the largest dimension. That is, the system was designed for the relatively large landscapes managed by most municipal, state and federal governments, and is often not appropriate for the small-scale landscapes studied by many academic biologists. Uncertainty about the scale at which users would apply the results was the prime reason for designing a modular system.

Although there is large variation in the sizes of RAPELD modules, most users use a standard 25 km² grid with 30 uniformly distributed plots for intensive studies in long-term ecological research sites located near major research institutions, and standard 5 km² (5 km × 1 km) modules with 10 uniformly distributed plots for RAP studies or long-term studies distributed over large areas (<http://ppbio.inpa.gov.br/inventarios/modular>). Uniformly distributed plots are 250 m long and the center line follows the altitudinal contours (<http://ppbio.inpa.gov.br/instalacao/parcelas>), a design that generally allows greater precision of models that relate biodiversity parameters to environmental variables. Plots for special strata, such as streams and riparian zones, are distributed in proportion to their occurrence in the landscape.

Most biologists specialize in a limited range of taxa (e.g. vascular plants) or processes (e.g. pollination), but decision makers have to take into account the needs of many different stakeholders, who may be interested in subjects as varied as the effect of large carnivores on domestic animals, bacterial metagenomics, ecosystem carbon storage and traditional uses of biodiversity. Reduction of these varied interests to a production-line mathematical model with limited inputs and outputs is usually not feasible, especially when a major uncertainty is whether we are addressing the right question (Haila et al. 2014). Therefore, investing monitoring in a limited number of questions, however important they may be at the moment, is not an efficient strategy.

A major difficulty, perhaps the major difficulty, with the interpretation of data collected in monitoring exercises is that the biologists have focused on their favorite group and not collected the data in such a way that it can be integrated with information generated on other biological groups and presumed environmental drivers. Different taxa provide different information about the distribution of biodiversity, and there is often heated discussion about the appropriate group to study (Magurran and McGill 2011). However, the sad reality is that we generally just base our decisions on convenience. There are few groups that have been surveyed over wide areas for which we are reasonably confident that most individuals have been correctly identified. These are usually only birds and vascular plants, though some groups of butterflies and mammals are reasonably well known in some areas. There is strong evidence that they are not sufficient to represent all biodiversity (Caro

2010), but they will continue to do so until we move out of our comfort zones within the internet cloud.

When we started our studies, discussing possible sampling designs only resulted in endless discussions as to which design optimized for a particular question was the “right” design. We found that the only way to obtain integrated data collection was to provide standardized infrastructure that could be used by most researchers to answer a wide variety of questions. Henle et al. (2006) present a European example of this approach, and Olsen et al. (2012) give several examples from the USA. Most researchers had not adequately budgeted for field infrastructure and were happy to use what was available. As this infrastructure had been designed to allow integration, data collection resulted in integrated studies almost as a side effect (Costa and Magnusson 2010, Magnusson et al. 2013).

There is a great difference between planning for an individual study of a limited range of organisms and planning a monitoring system for a wide range of taxa over very large areas. We did not appreciate this at the beginning, and it only became obvious to us as we saw what worked for a wide range of researchers over large areas, and what was mainly useful for specific studies. This dichotomy has been recognized by many researchers responsible for nationwide monitoring of biodiversity (e.g. Johnson 2012, Olsen et al. 2012), but has only recently been included in reviews of best monitoring practice from a statistical perspective (e.g. Buckland et al. 2011, Connolly and Dornelas 2011, Reynolds 2012), most of which had previously concentrated on idiosyncratic planning of projects with a common source of funding (e.g. Likens and Lindenmayer 2011). Below I will provide a short overview of key sources of uncertainty and how standardized field infrastructure can be used to help avoid or quantify uncertainty during the establishment and running of large-scale monitoring schemes.

Uncertainty about the stakeholders

Individual researchers tend to consider their study site to be primordially of interest in relation to their current research question. However, that piece of land may have a multitude of other values for the local people (Silvius et al. 2004). When planning where to install long-term research sites, we found that many different stakeholders were interested in the same site. Data generated might be used by international organizations, such as the International Long Term Ecological Research program and United Nations agencies, federal agencies, such as ministries of science and environment, regional bodies, such as State Government planning agencies, individuals operating regionally, such as university professors, park administrators and firms specialized in bioprospecting, and those interested in a small patch within a site, such as students, community groups and volunteers. In general they follow a political hierarchy (e.g. Magnusson et al. 2013: 58-59), but the categories do not always have clear boundaries. For example, individual volunteers may contribute to nation-wide projects, such as Christmas bird counts.

Stakeholder roles also depend on their position in the hierarchy. Multinational bodies try to influence decisions by changing national policies. Most funding agencies for long-term monitoring are national, but international organizations may provide short-term funding. Federal and state governmental agencies generally try to manipulate people's behavior through the legal system. Academics are involved in planning and analysis, and most of the long-term monitoring has to be done by local people or students. All of these categories have fuzzy boundaries, and the relative interest of these groups is likely to change depending on unpredictable factors, such as employment opportunities, market demand and global climate change. Nevertheless, a monitoring scheme has to take into account the different roles of stakeholders. Probably the most difficult aspect of developing the RAPELD system was ensuring that different actors in different levels of the political system would be satisfied with their role, and the roles of other actors (Magnusson et al. 2013).

Biological relationships with distance are not linear (Landeiro and Magnusson 2011, Magurran 2011, Rosenzweig et al. 2011). It is possible to scale up from local data to larger areas, but only if data are collected in spatially standardized arrays, and this may create conflicts with organizations and volunteers who collect the data (Turnhout and Boonman-Berson 2011). Relationships depend heavily on the sampling scale (Baccaro et al. 2012, Rosenzweig et al. 2011, Chisholm et al 2013). Evaluations of the effects of scale of sampling are generally difficult with idiosyncratic sampling, but can be achieved with a few sites with standardized infrastructure. Where coverage is inadequate, geostatistical techniques may help to define priorities for locations of new sampling sites (Lin et al. 2008). Modular designs allow flexibility in answering local questions, while permitting different stakeholders to adjust the system to their questions (Magnusson et al. 2013). Some RAPELD sites, such as that in the Virua National Park, have been used both for local studies of interest to park managers (e.g. Pontes et al. 2012) and included in cross-site comparisons for academic studies (e.g. Souza et al. 2012).

Uncertainty about who will monitor

Different stakeholders have different human and financial resources, but very few have the capacity to undertake detailed studies over large areas. Therefore, we needed a system that would allow integration of a large number of stakeholders with different technological tools at their disposal. Our infrastructure is suitable for use by local people with no formal education, and their participation is often vital because much biodiversity is hidden from the eyes of casual visitors (Magnusson et al. 2013).

Students are the main researchers in most RAPELD sites, but many of the surveys carried out in the modules around the Santo Antônio hydro-electric dam were undertaken by parataxonomists who had been trained in another state. We found that there was a trade-off in sources of uncertainty. Monitoring by students and volunteers

increases uncertainty as to the frequency and quality of monitoring. Relying on surveyors specifically contracted for the task, as was the case in Santo Antônio, increases uncertainty as to whether funding will be sufficiently reliable to meet labor-law requirements. By concentrating on field infrastructure, we were able to take advantage of different forms of financing for monitoring in different places and time periods.

Costs of monitoring could not be too great if different stakeholders with limited economic resources were to be involved, but the system had to house high-technology systems, such as eddy-flux towers, when available. While it is not possible to foresee all stakeholders, it is possible to provide field infrastructure that most will need, such as access trails and permanent plots.

Uncertainty about relations with remote sensing

Because of its complexity, all aspects of biodiversity cannot be measured directly and decisions are made based on surrogates, which are usually maps derived from remote-sensing data, but may be simply the representation of one biological group by another. For instance, vascular plants are often used to identify “habitats”, “ecoregions” or “ecosystems” that purportedly represent boundaries to the distribution of other organisms, such as insects, mammals or fish (e.g. Olson et al. 2001, Higgins et al. 2005). The technology is continually advancing, and it is not possible to predict what remote-sensing products will be available in the future, but the greatest sources of uncertainty at the moment relate to the relationships between surrogates and the target organisms in which we are interested (Magnusson 2004, Franklin 2009, Caro 2010).

Biologists generally stratify and collect only where they “know” that certain types of organisms occur (Henle et al. 2006). When they are forced to sample regularly or randomly because that is where the infrastructure is, they almost always discover that their preconceived ideas were wrong (Oliveira et al. 2008), including ideas about the relationships between species distributions and remote-sensing surrogates. Other researchers involved in country-wide programs have found a priori stratification to be problematical (e.g. Johnson 2012, McDonald 2012). Infrastructure that is not dedicated to particular groups reduces the risk that spatial sampling will be dedicated to a particular taxon. Standardized infrastructure in RAPELD plots has allowed validation of surrogates for which geographic information system (GIS) layers were available (e.g. Schietti et al. 2013).

We were initially uncertain about both the questions that stakeholders would want to answer and the remote-sensing technology that would become available. However, most political decisions are made on scales of tens to hundreds of linear kilometers, and few researchers have the resources to use remote-sensing tools with pixel sizes of a few meters. Therefore, we designed a system with relatively large sampling units (250 m long plots and 5 km long transects) that would allow the use of a wide variety of remote-sensing products available today, and will allow the use of even more in the future as products with smaller pixel sizes come on line.

Uncertainty about what to sample

Many monitoring programs have fixed targets, such as the Alberta Biodiversity Monitoring Institute (ABMI – Haughland et al. 2009), the Center for Tropical Forest Science (CTFS – Condit 1988) and Tropical Ecology, Assessment and Monitoring (TEAM – Martins et al. 2007). Standardization of a limited range of targets (e.g. vascular plants in CTFS plots) facilitates standardization, but leaves much biodiversity without coverage. Standardization for many groups (e.g. ABMI and TEAM) greatly increases the costs, and all standardization of targets reduces the range of stakeholders who will participate and the range of questions that can be answered. Therefore, we developed a system that permits essentially all elements of biodiversity to be monitored, but does not require that all targets are monitored in every site.

Integration can be obtained by associating individual sites with larger initiatives for particular targets. For instance, RAPELD plots are included in the RAINFOR (e.g. Emílio et al. 2013), GVID (e.g. Pezzini et al. 2012) and ATDN (Stropp et al. 2009, ter Steege 2013) vegetation networks. The same plots have been surveyed for taxa as diverse as mites (Franklin et al. 2013), ants (Souza et al. 2012), frogs (Menin et al. 2007) and birds (Bueno et al. 2012). There is a logical trade-off. Very strong standardization reduces coverage, but too little standardization makes wide-scale syntheses impossible. Discovering the most appropriate targets for any particular question is an on-going process, so making a design that is only appropriate for one biological group is not an optimal strategy.

Uncertainty about where was sampled

Most biologists now carry GPS devices, and geographic coordinates are the backbone of the Darwin-core system for digitalizing the information in biological collections. However, precise information about collection locations, or a single point representing the headquarters of park personnel in a large reserve are generally not sufficient to evaluate search effort or relate biological data to potential abiotic drivers. When we installed RAPELD modules in areas that had been intensively studied by other monitoring programs, we encountered many difficulties in avoiding disturbance to their plots, because the other programs did not have precise coordinates delimitating their field infrastructure.

RAPELD modules provide researchers with extremely detailed information on the location of trails, plots and large sessile organisms, such as trees. As RAPELD plots are long and thin, it is easy for researchers to locate their organisms quickly using only a compass and a measuring tape. All trails are marked at 50 m intervals (100 m intervals in some older sites), so researchers and local assistants can record relatively precise locations even when they do not have GPS equipment. This has been especially important for the use of RAPELD in environmental-impact studies, because reduction in area occupied is often a more sensitive measure of impact than attempts to estimate absolute numbers of organisms by mark-recapture methods (See “Uncertainty about detection” below).

Uncertainty about relationships

Different aspects of biodiversity and the environment are usually studied by different researchers, and very often it is difficult to see what these researchers have in common. “Integrated” projects usually involve extensive discussion about how the funds will be divided and the general locations of study sites. However, when the time comes for analysis, it is usually impossible to integrate the different studies that were done on different temporal and spatial scales, even though all were contained within the same geographic envelope. Often, researchers from different disciplines have completely different concepts of what represents replication and independence of observations.

By installing a system of transects, and especially plots, that can be used by a wide range of disciplines, we were able to integrate many studies that previously had been considered too disparate for interdisciplinary studies (see also Henle et al. 2006). Almost all studies that have come out of RAPELD systems have integrated data from a variety of disciplines, but have generally focused on a limited range of biological taxa. However, the RAPELD spatial standardization and emphasis on data storage and availability have allowed integrated studies of concordance in the landscape distribution of different taxa (e.g. Landeiro et al. 2012).

Narrow plots that follow altitudinal contours have less internal variability in environmental predictor variables than conventional square plots, or long thin plots that are not oriented along contour lines (Henle et al. 2006, Castilho et al. 2010). This allows much more precise determination of relationships with predictors than possible with standard plots. As important as reducing uncertainty, is that this reduces costs, and allows detection of changes within a shorter period. For instance, Castilho et al. (2010) were able to show significant relationships between biomass accumulation and environmental predictors with 2-yr intervals between tree surveys in a RAPELD grid, at a fraction of the cost of the implementation of a single large plot, even though the differences would have been within the measurement error for a single large plot.

Most studies in RAPELD plots have investigated the relationships between topographic, soil or hydrological variables and organisms (e.g. Menin et al. 2007, Bueno et al. 2012, Baccaro et al. 2013, Emílio et al. 2013, Schietti et al. 2013), but some studies have investigated relationships among different groups of organisms (e.g. Baccaro et al. 2012, Landeiro et al. 2012) and studies are starting to investigate how relationships change over time (e.g. Espírito-Santo et al. 2009, Castilho et al. 2010). However, evaluation of geographic variation in these relationships based on RAPELD spatial designs replicated in different regions has been undertaken for few groups (e.g. Souza et al. 2012, Zuquim et al. 2012).

Uncertainties that can be reduced by quality control

Uncertainty about identifications

Field work is increasingly being considered unfashionable (Magnusson 1994) and many believe that we can resolve all the problems associated with biodiversity by min-

ing digital information on collections. However, the quality of identifications is extremely varied. An internet search for photographs of a given species will often result in photographs of organisms from different genera or even phyla. For bio-diverse groups, such as arthropods, the species may not have been described. Advances in genetic techniques may alleviate these problems for collected specimens in the future, but many surveys, especially of endangered taxa, are based on sightings.

Different observers can result in different diversities, and different levels of biotic complementarity, even for comparisons of the same site. Therefore, we have invested heavily in surveyor mobility, so that different researchers can exchange experiences and compare identifications. Field workshops are much more efficient than learning by reading, and field courses are a large part of our investment <<http://ppbio.inpa.gov.br/extensao>>. In any case, as many voucher specimens and photographs should be taken as are financially and ethically feasible. Passive sampling by traps may reduce observer biases, and some taxa can only be efficiently sampled with traps. However, passive sampling is often inefficient in comparison to active sampling (e.g. Ellison et al. 2007), and passive sampling usually does not allow precise evaluation of the area sampled by the device, invalidating estimates of the number of species in the area of interest using rarefaction.

Marked plants (live herbariums) may allow re-evaluation of identifications in the future. Production of printed and internet field guides helps maintain stability of identifications across sites, and within the same site through time. For the first RAPELD site, we produced guides to frogs, lizards, the predominant understory angiosperms and ferns (Costa et al. 2008, Lima et al. 2008, Vitt et al. 2008, Zuquim et al. 2008). Guides to snakes, ants and fungal fruiting bodies are in production. Video footage and sonograms help identification of many taxa <<http://ppbio.inpa.gov.br/sapoteca/paginainicial>>. Where possible, genetic material should be collected and stored, even if resources are not currently available for analyses. The laboratory costs for genetic analyses are dropping precipitately, and exotic and expensive analyses today will be routine in the future. Other techniques, such as near-infrared spectroscopy (NIRS), offer even cheaper solutions (Foley et al. 1998, Durgante et al. 2013). However, if the material is not collected now, we may lose our bench marks.

Uncertainty about detectability

Conservation decisions are made based on the distribution of taxa, but distribution is defined as much by the area that a species does not occur as by where it occurs. False absences may lead to bad scientific decisions (Yoccoz et al. 2001, Buckland et al. 2011), and those decisions may lead to the waste of limited conservation resources. There are many methods for correcting for the detectability of individuals (Williams et al. 2002, Buckland et al. 2011), but they are usually too expensive to be applied in general surveys. In contrast, evaluation of the detectability of species may allow much better estimates of the proportion of the landscape occupied by a given taxon (MacKenzie et al. 2002).

Estimates of species detectability and occupancy generally require repeated surveys of sampling units, though in some cases space can be substituted for time. To be able to

use those techniques, it is important that detailed information is available about where organisms were collected, and the effort expended to detect them. This is relatively easy to do with the spatially standardized sampling units used in the RAPELD system. Precise coordinates are often available for specimens collected in conventional surveys, but researchers who do not collect in spatially standardized units usually do not report sampling effort, especially if no specimen was collected. Spatially standardized units have allowed the evaluation of occupancy in RAPELD modules, and will allow long-term changes in occupancy to be evaluated. Even in the case that the researcher is confident that they record all the species within a sampling unit (a rare occurrence in the field, but common in researcher imaginations), quantifying detectability greatly increases the confidence that other researchers and managers will have in the results.

Uncertainty about complementarity

Complementarity is a core concept in systematic conservation planning (Jost et al. 2011, Kukkala and Moilanen 2013). However, most of the software used assumes that there is complete knowledge of the distribution of species, and this is not the case for many bio-diverse taxa in bio-diverse regions (see sections on detectability and distributions above). Studies by Ana Albernaz in the Alter do Chaó region showed that, when these assumptions are not met, results reflect more the distribution of sampling than the distribution of the biota (Magnusson et al. 2013).

There are other options for planning, such as the use of complementarity of species assemblages based on multivariate ordination techniques (e.g. Reyers et al. 2002, Ilg et al. 2012) and selection of sites for additional sampling when initial sampling is inadequate (e.g. Lin et al. 2008). However, spatially standardized sampling is generally the easiest option to evaluate the assumptions of analyses, and standardized sampling has allowed evaluation of the effects of scale and position of sampling units in RAPELD plots (Franklin et al. 2013) and elsewhere (Chisholm et al. 2013).

Uncertainty about data quality and management

For monitoring, data is generally a more important product than scientific publications (Costello et al. 2013, Piwowar 2013). There are many schemes for storing and making data available, often with automatic upload by individual researchers. However, just having the system available does not mean that it will be used, leading to the “empty archive” syndrome (Nelson 2009). Also, data with errors may be worse than no data at all. We were unable to achieve adequate data quality just by offering information-technology resources; we had to have a human in the system (Pezzini et al. 2012, Magnusson et al. 2013). Other organizations have come to the same conclusion (Billick 2010). The spatially standardized system facilitated the production of data forms, but the flexibility in the taxa and collecting techniques added to the complexity of the process.

We were initially uncertain as to the best way to make data available. There are many database programs available, and information-technology specialists are always willing to develop another one. However, use of a single database for all the data is not viable for diverse monitoring data (Hale 1999), and it is more efficient to use a data repository that can be used to populate different databases with different purposes (Reichman et al. 2011). In this case, searches are undertaken on the metadata rather than the data themselves (Evans and Foster 2011). In the end, we found that the technology available far exceeds the ability of researchers to use it.

The information-technology revolution is recent, and none of our major field researchers had formal training in data management. Worse still, most university programs still do not offer specific courses in the principles of data management to biologists, though some do offer courses in the use of specific database programs. Data management is not easy or intuitive, and it is a critical phase in research that can effectively nullify all the planning that has gone into data collection. In the end, we found that the major uncertainties were related to whether (1) researchers would be motivated to make their data available, (2) whether researchers had already lost critical data in the field, (3) whether researchers had sufficient training to effectively deposit data, and (4) whether we could find the resources to undertake the capacity building and data verification necessary. Every major project should have a full time data manager responsible for training and data screening, but few do.

We adopted the Ecological Metadata Language (EML) used in the Metacat system (Fegraus et al. 2005) by the International Long-Term Ecological Research (ILTER) sites. However, we found that the generic coordinates required by that system, which basically just locate the study site, were not sufficient for most ecological analyses, especially of local landscapes. As the sampling sites are standardized, it was relatively easy for us to annex accessory metadata tables <http://ppbio.inpa.gov.br/sites/default/files/repositorio_PPBio_maio_2012.rar> to the EML metadata so that researchers could easily record the detailed information that will make their data useful to the broadest range of researchers in the future.

Conclusions

I have covered only a few of the sources of uncertainty that a field monitoring scheme has to deal with, but they illustrate the complexity of the problem. We are monitoring because we are uncertain, so uncertainty has to be a central issue in any monitoring scheme. However, biodiversity monitoring is more complex than monitoring physical phenomena, such as weather, because there are so many definitions of biodiversity, and it has different values for different segments of society (see Haila et al. 2014).

When we first implemented RAPELD, we opted for a hierarchical design, with regular sampling at the local level, but sampling sites limited by logistical considerations over larger areas. Although there was little theoretical support for it at the time, it has been useful to respond to a wide range of questions of interest to decision makers (Magnusson et al. 2013),

it is similar to designs adopted by other researchers faced with sampling biodiversity over large areas (e.g. McDonald 2012, Olsen et al. 2012), and it is a design that has since been approved by experienced modelers for continent-wide monitoring (e.g. Franklin 2009, Johnson 2012, Reynolds 2012). Because the design is modular, it can be adapted to a wide range of situations. At first, we envisaged that most sites would have 5 km × 5 m grids, but the most favored module today is 5 km × 1 km. We were uncertain about everything and adopted a learning-by-doing approach, while always trying to maximize comparability with data collected previously. We are more surprised at how little we have had to modify the original design than by the changes we have implemented (Magnusson et al. 2013).

We were even uncertain about the questions that researchers will want to answer in the future using the data we are collecting today. Therefore, rather than adopting a taxon- or question-oriented approach, we focused on a spatial design that allows flexibility in questions and taxa studied, while allowing the landscape-geographical approaches required by conservation biology. It is a compromise, but a necessary one. RAPELD sites are being used to answer many specific questions, so the trade-off is generally not between monitoring and answering specific questions. It is between investing in field infrastructure and planning now, rather than in short-term studies that are good for researcher curricula, but that contribute little to long-term, wide-scale conservation planning. At the moment, many citizen monitoring programs, combined with good data availability, are contributing more to our understanding of global phenomena, such as climate change, than are more scientific programs (Schmeller et al. 2009), despite criticisms from academics (e.g. Ferraz et al. 2008). We professional scientists need to learn from the amateurs, and incorporate planning, standardization, and data availability, while maintaining the flexibility that uncertainty demands.

Acknowledgments

The ideas expressed in this paper were developed by too many people to cite here, but all are participants in the Instituto Nacional de Ciência Tecnologia - Centro de Estudos Integrados da Biodiversidade Amazônica (INCT-CENBAM), and the Brazilian Program for Biodiversity Research (PPBio). I thank Klaus Henle for the opportunity to participate in the workshop and the stimulus for the important interactions through EU Bon (EU FP project grant number 308454).

References

- Anderson DR (2008) *Model Based Inference in the Life Sciences*. Springer, New York, 184 pp. ISBN 978-0-387-74075-1
- Baccaro FB, De Souza JLP, Franklin E, Landeiro VL, Magnusson WE (2012) Limited effects of dominant ants on assemblage species richness in three Amazon forests. *Ecological Entomology* 37: 1–12. doi: 10.1111/j.1365-2311.2011.01326.x

- Baccaro FB, Rocha IF, Aguila EG, Schiatti J, Emílio T, Pinto JLPV, Lima AP, Magnusson WE (2013) Changes in ant functional groups are correlated with water-table level in an Amazonian terra firme forest. *Biotropica* 45: 755–763. doi: 10.1111/btp.12055
- Billick I (2010) Managing place-based data. In: Billick I, Price MV (Eds) *The Ecology of Place*. University of Chicago Press, Chicago, IL, 382–402. doi: 10.7208/chicago/9780226050447.001.0001
- Buckland ST, Studeny AC, Magurran AE, Newson SE (2011) Biodiversity monitoring: the relevance of detectability. In: Magurran AE, McGill BJ (2011) *Biological Diversity*. Oxford University Press, Oxford, 25–36.
- Bueno AS, Bruno RS, Pimentel TP, Sanaiotti TM, Magnusson WE (2012) The width of riparian habitats for understory birds in an Amazonian forest. *Ecological Applications* 22: 722–734. doi: 10.1890/11-0789.1
- Caley P, Hone J (2002) Estimating the force of infection: *Mycobacterium bovis* infection in feral ferrets *Mustela furo* in New Zealand. *Journal of Animal Ecology* 71: 44–54. doi: 10.1046/j.0021-8790.2001.00573.x
- Caro T (2010) *Conservation by Proxy: Indicator, Umbrella, Keystone, Flagship, and Other Surrogate Species*. Island Press, Washington, 400 pp.
- Castilho CV, Magnusson WE, Araújo RNO, Luizaõ FJ (2010) Short-term temporal changes in tree live biomass in a Central Amazonian forest, Brazil. *Biotropica* 42: 95–103. doi: 10.1111/j.1744-7429.2009.00543.x
- Chisholm RA, Muller-Landau HC, Rahman KA, Bebbler DP, Bin Y, Bohlman SA, Bourg NA, Brinks J, Bunyavejchewin S, Butt N, Cao H, Cao M, Cardenas D, Chang L-W, Chiang J-M, Chuyong G, Condit R, Dattaraja HS, Davies S, Duque A, Fletcher C, Gunatilleke N, Gunatilleke S, Hao Z, Harrison RD, Howe R, Hsieh C-F, Hubbell SP, Itoh A, Kenfack D, Kiratiprayoon S, Larson AJ, Lian J, Lin D, Liu H, Lutz JA, Ma K, Malhi Y, McMahon S, McShea W, Meegaskumbura M, Razman SM, Morecroft MD, Nytech CJ, Oliveira A, Parker GG, Pulla S, Punchi-Manage R, Romero-Saltos H, Sang W, Schurman J, Su1S-H, SukumarR, SunI-F, Suresh HS, Tan S, Thomas D, Thomas S, Thompson J, Valencia R, Wolf A, Yap S, Ye W, Yuan Z, Zimmerman JK (2013) Scale-dependent relationships between tree species richness and ecosystem function in forests. *Journal of Ecology* 101: 1214–1224. doi: 10.1111/1365-2745.12132
- Condit R (1988) *Tropical Forest Census Plots*. Springer, New York, NY, 221 pp.
- Connolly SR, Dornelas M (2011) Fitting and empirical evaluation of models for species abundance distributions. In: Magurran AE, McGill BJ (2011) *Biological Diversity*. Oxford University Press, Oxford, UK, 123–140.
- Costa FRC, Magnusson WE (2010) The need for large-scale, integrated studies of biodiversity – the experience of the program for biodiversity research in Brazilian Amazônia. *Natureza & Conservação* 8: 1–5. doi: 10.4322/natcon.00801001
- Costa FRC, Espinelli FP, Figueiredo FO (2008) *Guide to the Marantaceae of Reserva Ducke & Reserva Biológica Uatumã*. Áttema Design Editorial, Manaus, AM.
- Costello MJ, Michener WK, Gahegan M, Zhang Z-Q, Bourne PE (2013) Biodiversity data should be published, cited, and peer reviewed. *Trends in Ecology & Evolution* 28: 454–461. doi: 10.1016/j.tree.2013.05.002

- Durgante FM, Higuchi N, Vicentini A, Almeida A (2013) Species spectral signature: Discriminating closely related plant species in the Amazon with near-infrared leaf-spectroscopy. *Forest Ecology and Management* 291: 240–248. doi: 10.1016/j.foreco.2012.10.045
- Ellison AM, Record S, Arguello A, Gotelli NJ (2007) Rapid inventory of the ant assemblage in a temperate hardwood forest: species composition and assessment of sampling methods. *Environmental Entomology* 36: 766–775. doi: 10.1603/0046-225X(2007)36[766:RIOTAA]2.0.CO;2
- Emilio T, Quesada CA, Costa FRC, Magnusson WE, Schiatti J, Feldpausch TR, Brienen RJW, Baker TR, Chave J, Álvarez E, Araújo A, Bánki O, Castilho CV, Honorio C, Killeen TJ, Malhi Y, Mendoza EMO, Monteagudo A, Neill D, Parada GA, Pe-a-Cruz A, Ramirez-Angulo H, Schwarz M, Silveira M, ter Steege H, Terborgh JW, Thomas R, Torres-Lezama A, Vilanova E, Phillips OL (2013) Soil physical conditions limit palm and tree basal area in Amazonian forests. *Plant Ecology and Diversity* 7(1/2): 215–229. doi: 10.1080/17550874.2013.772257
- Espírito-Santo HVM, Magnusson WE, Zuanon J, Mendonça FP, Landeiro VL (2009) Seasonal variation in the composition of fish assemblages in small Amazonian streams: evidence for predictable changes. *Freshwater Biology* 54: 536–548. doi: 10.1111/j.1365-2427.2008.02129.x
- Evans JÁ, Foster JG (2011) Metaknowledge. *Science* 331: 721–725. doi: 10.1126/science.1201765
- Fegraus EH, Andelman S, Jones MB, Schildhauer M (2005) Maximizing the value of ecological data with structured metadata: An introduction to ecological metadata language (EML) and principles for metadata creation. *Bulletin of the Ecological Society of America* 86: 158–168. doi: 10.1890/0012-9623(2005)86[158:MTVOED]2.0.CO;2
- Ferraz G, Marinelli CE, Lovejoy TE (2008) Biological monitoring in the Amazon: Recent progress and future needs. *Biotropica* 40: 7–10.
- Foley WJ, McIlwee A, Lawler I, Aragones L, Woolnough AP, Berding N (1998) Ecological applications of near infrared reflectance spectroscopy - a tool for rapid, cost-effective prediction of composition of plant and animal tissues and aspects of animal performance. *Oecologia* 116: 293–305. doi: 10.1007/s004420050591
- Franklin J (2009) *Mapping Species Distributions*. Cambridge University Press, Cambridge, 336 pp.
- Franklin E, Moraes J, Landeiro VL, Souza JLP, Pequeno PACL, Magnusson WE, Morais JW (2013) Geographic position of sample grid and removal of uncommon species affect multivariate analyses of diverse assemblages: The case of oribatid mites (Acari: Oribatida). *Ecological Indicators* 34: 172–180. doi: 10.1016/j.ecolind.2013.04.024
- Gitzen RA, Millsbaugh JJ, Cooper AB, Licht DS (Eds) (2012) *Design and Analysis of Long-term Ecological Monitoring Studies*. Cambridge University Press, Cambridge, UK, 590 pp. doi: 10.1017/CBO9781139022422
- Haila Y, Henle K, Apostolopoulou E, Joanna Cent J, Framstad E, Görg C, Jax K, Klenke R, Magnusson WE, Matsinos Y, Müller B, Paloniemi R, Pantis J, Rauschmayer F, Ring I, Settele J, Similä J, Touloumis K, Tzanopoulos J, Pe'er G (2014) Confronting and Coping with Uncertainty in Biodiversity Research and Praxis. *Nature Conservation* 8: 45–75. doi: 10.3897/natureconservation.8.5942

- Hale SS (1999) How To Manage Data Badly (Part 1). *Bulletin of the Ecological Society of America* 80: 265–268.
- Haughland DL, Hero J-M, Schieck J, Castley JG, Boutin S, Sóllymos P, Lawson BE, Holloway G, Magnusson WE (2009) Planning forwards: biodiversity research and monitoring systems for better management. *Trends in Ecology and Evolution* 25: 199–200. doi: 10.1016/j.tree.2009.11.005
- Henle K, Dziöck F, Foeckler F, Follner K, Hüsing V, Hettrich A, Rink M, Stab S, Scholz M (2006) Study Design for assessing species environment relationships and developing indicator systems for ecological changes in floodplains – The approach of the RIVA Project. *International Review Hydrobiology* 91: 292–313. doi: 10.1002/iroh.200610886
- Higgins JV, Bryer MT, Khoury ML, Fitzhugh TW (2005) Freshwater classification approach for biodiversity conservation planning. *Conservation Biology* 19: 432–445. doi: 10.1111/j.1523-1739.2005.00504.x
- Ilg C, Foeckler F, Deichner O, Henle K (2012) Hydrological gradient and species traits explain gastropod diversity in floodplain grasslands. *River Research and Applications* 28: 1620–1629. doi: 10.1002/rra.1552
- Johnson DH (2012) Monitoring that matters. In: Gitzen RA, Millsbaugh JJ, Cooper AB, Licht DS (Eds) *Design and Analysis of Long-term Ecological Monitoring Studies*. Cambridge University Press, Cambridge, 54–73. doi: 10.1017/CBO9781139022422.006
- Jost L, Chao A, Chazdon RL (2011) Compositional similarity and β (beta) diversity. In: Magurran AE, McGill BJ (Eds) *Biological Diversity*. Oxford University Press, Oxford, 66–84.
- Kukkala AS, Moilanen A (2013) Core concepts of spatial prioritization in systematic conservation planning. *Biological Reviews* 88: 443–464. doi: 10.1111/brv.12008
- Landeiro VL, Magnusson WE (2011) The geometry of spatial analyses: implications for conservation biologists. *Natureza & Conservação* 9(1): 7–20. doi: 10.4322/natcon.2011.002
- Landeiro VL, Bini LM, Costa FRC, Franklin E, Nogueira A, Souza JLP, Moraes J, Magnusson WE (2012) How far can we go in simplifying biomonitoring assessments? An integrated analysis of taxonomic surrogacy, taxonomic sufficiency and numerical resolution in a megadiverse region. *Ecological Indicators* 23: 366–373. doi: 10.1016/j.ecolind.2012.04.023
- Likens GE, Lindenmayer DB (2011) A strategic plan for an Australian Long-Term Environmental Monitoring Network. *Austral Ecology* 36: 1–8. doi: 10.1111/j.1442-9993.2010.02179.x
- Lima AP, Magnusson WE, Menin M, Erdtmann LK, Rodrigues DJ, Keller C, Hödl W (2008) *Guide to the Frogs of Reserva Adolpho Ducke: Central Amazonia*. Áttema Design Editorial, Manaus.
- Lin Y-P, Yeh M-S, Deng D-P, Wang Y-C (2008). Geostatistical approaches and optimal additional sampling schemes for spatial patterns and future sampling of bird diversity. *Global Ecology, Biogeography* 17: 175–188. doi: 10.1111/j.1466-8238.2007.00352.x
- MacKenzie DI, Nichols JD, Lachman GB, Droege S, Royle JA, Langtimm CA (2002) Estimating site occupancy rates when detection probabilities are less than one. *Ecology* 83: 2248–2255. doi: 10.1890/0012-9658(2002)083[2248:ESORWD]2.0.CO;2

- Magnusson WE (1994) The role of wildlife management in the maintenance of biodiversity. In: Majumdar SK, Brenner FJ, Lovich JE, Schalles JF, Miller EW (Eds) *Biological Diversity: Problems and Challenges*. Pennsylvania Academy of Science, Easton (Pennsylvania), 209–220.
- Magnusson WE (2004) Ecoregion as a pragmatic tool. *Conservation Biology* 18: 4–5. doi: 10.1111/j.1523-1739.2004.t01-1-00L13.x
- Magnusson WE, Lima AP, Luizão R, Luizão F, Costa FRC, Castilho CV, Kinupp VF (2005) RAPELD: a modification of the Gentry method for biodiversity surveys in long-term ecological research sites. *Biota Neotropica* 5: 19–24. doi: 10.1590/S1676-06032005000300002
- Magnusson W, Braga-Neto R, Pezzini F, Baccaro F, Bergallo H, Penha J, Rodrigues D, Verdade L, Lima A, Albernaz, A, Hero J-M, Lawson B, Castilho C, Drucker D, Franklin E, Mendonça F, Costa F, Galdino G, Castley G, Zuanon J, Vale J, Santos J, Luizão R, Cintra R, Barbosa R, Lisboa A, Koblitz R, Cunha, C, Mendes Pontes A (2013) Biodiversity and Integrated Environmental Monitoring. *Áttema Editorial*, Manaus. <http://ppbio.inpa.gov.br/sites/default/files/Biodiversidade%20e%20monitoramento%20ambiental%20integrado.pdf>
- Magurran AE (2011) Measuring biological diversity in time (and space). In: Magurran AE, McGill BJ (Eds) *Biological Diversity*. Oxford University Press, Oxford, 85–94.
- Magurran AE, McGill BJ (2011) *Biological Diversity*. Oxford University Press, Oxford, 368 pp.
- McDonald T (2012) Spatial sampling designs for long-term ecological monitoring. In: Gitzen RA, Millspaugh JJ, Cooper AB, Licht DS (Eds) *Design and Analysis of Long-term Ecological Monitoring Studies*. Cambridge University Press, Cambridge, 101–125. doi: 10.1017/CBO9781139022422.009
- Menin M, Lima AP, Magnusson WE, Waldez F (2007) Topographic and edaphic effects on the distribution of terrestrially reproducing anurans in Central Amazonia: mesoscale spatial patterns. *Journal of Tropical Ecology* 23: 539–547. doi: 10.1017/S0266467407004269
- Nelson B (2009) Data sharing: Empty archives. *Nature* 461: 160–163. doi: 10.1038/461160a
- Olsen AR, Kincaid TM, Payton Q (2012) Spatially balanced survey designs for natural resources. In: Gitzen RA, Millspaugh JJ, Cooper AB, Licht DS (Eds) *Design and Analysis of Long-term Ecological Monitoring Studies*. Cambridge University Press, Cambridge, 126–150. doi: 10.1017/CBO9781139022422.010
- Oliveira ML, Baccaro FB, Braga-Neto R, Magnusson WE (2008) Reserva Ducke: A Biodiversidade Através de uma Grade. *Áttema Design Editorial*, Manaus.
- Pezzini FF, Melo PHA, Oliveira DMS, Amorim RX, Figueiredo FOG, Drucker DP, Rodrigues FRO, Zuquim G, Emilio T, Costa FRC, Magnusson WE, Sampaio AF, Lima AP, Garcia ARM, Manzatto AG, Nogueira A, Costa CP, Barbosa CEA, Bernardes C, Castilho CV, Cunha CN, Freitas CG, Cavalcante CO, Brandão DO, Rodrigues DJ, Santos ECPR, Baccaro FB, Ishida FY, Carvalho FA, Moulatlet GM, Guillaumet J-LB, Pinto JLPV, Schiatti J, Vale JD, Belger L, Verdade LM, Pansonato MP, Nascimento MT, Santos MCL, Cunha MS, Arruda R, Barbosa RI, Romero RL, Pansini S, Pimentel TP (2012) The Brazilian Program for Biodiversity Research (PPBio) information System. *Biodiversity, Ecology* 4: 265–274.
- Pontes ARM, Paula MD, Magnusson WE (2012) Low primate density and abundance in northern Amazonia and its implications for conservation. *Biotropica* 44: 834–839. doi: 10.1111/j.1744-7429.2012.00873.x

- Reichman OJ, Jones MB, Schildhauer MP (2011) Challenges and opportunities of open data in ecology. *Science* 331: 703–705. doi: 10.1126/science.1197962
- Reyers B, Fairbanks DHK1, Wessels KJ, van Jaarsveld AS (2002) A multicriteria approach to reserve selection: addressing long-term biodiversity maintenance. *Biodiversity and Conservation* 11: 769–793. doi: 10.1023/A:1015315405292
- Reynolds JH (2012) An overview of statistical considerations in long-term monitoring. In: Gitzen RA, Millsaugh JJ, Cooper AB, Licht DS (Eds) *Design and Analysis of Long-term Ecological Monitoring Studies*. Cambridge University Press, Cambridge, 23–53. doi: 10.1017/CBO9781139022422.005
- Rosenzweig ML, Donoghue J, Li YM, Yuan C (2011) Estimating species density. In: Magurran AE, McGill BJ (Eds) *Biological Diversity*. Oxford University Press, Oxford, 276–288.
- Schmeller DS, Henry P-Y, Julliard R, Gruber B, Clobert J, Dziok F, Lengyel S, Nowicki P, D’eri E, Budrys E, Kull T, Tali K, Bauch B, Settele J, van Swaay C, Kobler A, Babij V, Pastergiadou E, Henle K (2009) Advantages of Volunteer-Based Biodiversity Monitoring in Europe. *Conservation Biology* 23: 307–316. doi: 10.1111/j.1523-1739.2008.01125.x
- Schiatti J, Emílio T, Renno CD, Drucker DP, Costa FRC, Nogueira A, Baccaro FB, Figueiredo F, Castilho CV, Kinupp V, Guillaumet J-L, Garcia ARM, Lima AP, Magnusson WE (2013) Vertical distance to drainage drives floristic composition changes in an Amazonian forest. *Plant Ecology, Diversity*. doi: 10.1080/17550874.2013.783642
- Silvius, KM, Bodmer RE, Fragoso, JMV (2004) *People in Nature: Wildlife Conservation in South and Central America*. Columbia University Press, New York, 464 pp.
- Snieidovich M (2014) Response to Burgman and Regan: the elephant in the rhetoric on info-gap decision theory. *Ecological Applications* 24: 229–233. doi: 10.1890/13-1096.1
- Souza JLP, Baccaro FB, Landeiro VL, Franklin E, Magnusson WE (2012) Trade-offs between complementarity and redundancy in the use of different sampling techniques for ground-dwelling ant assemblages. *Applied Soil Ecology* 56: 63–73. doi: 10.1016/j.apsoil.2012.01.004
- Stropp J, Ter Steege H, Malhi Y, ATDN, RAINFOR (2009) Disentangling regional and local tree diversity in the Amazon. *Ecography* 32: 46–54. doi: 10.1111/j.1600-0587.2009.05811.x
- ter Steege H, Pitman NCA, Sabatier D, Baraloto C, Salomão RP, Guevara JE, Phillips OL, Castilho CV, Magnusson WE, Molino J-F, Monteagudo A, Vargas PN, Montero JC, Feldpausch TR, Coronado HEN, Killeen TJ, Mostacedo B, Vasquez R, Assis RL, Terborgh J, Wittmann F, Andrade A, Laurance WF, Laurance SGW, Marimon BS, Marimon B-H, Vieira ICG, Amaral IL, Brienen R, Castellanos H, López DC, Duivenvoorden JF, Mogollón HF, Matos FDA, Dávila N, García-Villacorta R, Diaz PRS, Costa F, Emilio T, Levis C, Schiatti J, Souza P, Alonso A, Dallmeier F, Montoya AJD, Piedade MTF, Araujo-Murakami A, Arroyo L, Gribel R, Fine PVA, Peres CA, Toledo M, 39 Aymard C GA, Baker TA, Cerón C, Engel J, Henkel TW, Maas P, Petronelli P, Stropp J, Zartman CE, Daly D, Neill D, Silveira M, Paredes MR, Chave J, Lima Filho DA, Jørgensen PM, 51 Fuentes A, Schöngart J, Valverde FC, Di Fiore A, Jimenez EM, Mora MCP, Phillips JF, Rivas G, van Andel TR, von Hildebrand P, Hoffman B, Zent EL, Malhi Y, Prieto A, Rudas A, Ruschell AR, Silva N, Vos V, Zent S, Oliveira AA, Schutz AC, Gonzales T, Nascimento MT, Ramirez-Ángulo H, Sierra R, Tirado M, Medina MNU, van der Heijden G,

- Vela CIA, Torre EV, Vriesendorp C, Wang O, Young KR, Baider C, Balslev H, Ferreira C, Mesones I, Torres-Lezama A, Giraldo LEU, Zagrt R, Alexiades MN, Hernandez L, Huamantupa-Chuquimaco I, Milliken W, Cuenca WP, Pauletto D, Sandoval EV, Gamarra LV, Dexter KG, Feeley K, Lopez-Gonzalez G, Silman MR (2013) Hyperdominance in the Amazonian tree flora. *Science* 342: 325. doi: 10.1126/science.1243092
- Thornton JT, Cope JM, Patrick WS (2014) Assessing the quality of life history information in publicly available databases. *Ecological Applications* 24: 217–226. doi: 10.1890/12-1855.1
- Turnhout E, Boonman-Berson S (2011) Databases, scaling practices and globalization of biodiversity. *Ecology and Society* 16: 1–35. <http://www.ecologyandsociety.org/vol16/iss1/art35/>
- Vitt LJ, Magnusson WE, Avila-Pires TCS, Lima AP (2008) Guide to the Lizards of Reserva Adolpho Ducke, Central Amazonia. Áttema Design Editorial, Manaus.
- Williams BK, Nichols J, Conroy M (2002) Analysis and Management of Animal Populations. Academic Press, London, 817 pp.
- Yoccoz NG, Nichols JD, Boulinier T (2001) Monitoring of biological diversity in space and time. *Trends in Ecology, Evolution* 16: 446–453. doi: 10.1016/S0169-5347(01)02205-4
- Zuquim G, Costa FRC, Prado J, Tuomisto H (2008) Guide to the Ferns and Lycophytes of REBIO Uatumã, Central Amazonia. Áttema Design Editorial, Manaus.
- Zuquim G, Tuomisto H, Costa FRC, Prado J, Magnusson WE, Pimentel P, Braga-Neto R, Figueiredo FOG (2012) Broad-scale distribution of ferns and lycophytes along environmental gradients in central and northern Amazonia, Brazil. *Biotropica* 44: 752–762. doi: 10.1111/j.1744-7429.2012.00880.x